

Supplementary Table 1. Medication of the patients according to level of serum NT-proBNP and CgA

Variables	All (n = 272)	Low NT- proBNP (n = 146)	High NT- proBNP (n=126)	<i>P</i>	Low CgA (n = 166)	High CgA (n=106)	<i>P</i>
Medications before the admission							
Beta blockers	73 (26.8%)	41 (28.1%)	32 (25.4%)	0.718	44 (26.5%)	29 (27.4%)	0.988
ACE inhibitors	36 (13.2%)	20 (13.7%)	16 (12.7%)	0.950	26 (15.7%)	10 (9.4%)	0.195
Angiotensin receptor blockers	39 (14.3%)	21 (14.4%)	18 (14.3%)	1.000	27 (16.3%)	12 (11.3%)	0.338
MRA	36 (13.2%)	16 (11.0%)	20 (15.9%)	0.311	23 (13.9%)	13 (12.3%)	0.846
Loop diuretics	86 (31.6%)	38 (26.0%)	48 (38.1%)	0.045	44 (26.5%)	42 (39.6%)	0.033
Discharge medications							
Beta blockers	202 (76.2%)	112 (77.8%)	90 (74.4%)	0.615	119 (72.6%)	83 (82.2%)	0.102
ACE inhibitors	83 (31.4%)	52 (36.1%)	31 (25.8%)	0.097	63 (38.7%)	20 (19.8%)	0.002
ARB	67 (25.5%)	41 (28.7%)	26 (21.7%)	0.247	42 (25.9%)	25 (24.8%)	0.947
ARNI	12 (4.5%)	4 (2.8%)	8 (6.7%)	0.225	9 (5.5%)	3 (3.0%)	0.507
Ivabradine	11 (4.2%)	4 (2.8%)	7 (5.8%)	0.353	10 (6.1%)	1 (1.0%)	0.086

MRA	90 (34.1%)	34 (23.6%)	56 (46.7%)	0.000	56 (34.4%)	34 (33.7%)	1.000
Loop diuretics	191 (72.3%)	86 (59.7%)	105 (87.5%)	0.000	106 (65.0%)	85 (84.2%)	0.001
